PhytoGreen®-HiPhos

PK FERTILISER SOLUTION with Magnesium 31-5 (5,6)

Composition:

30,6 % P_2O_5 (440 g/l water soluble phosphate)
5,1 % K_2O (74 g/l water soluble potassium oxide)
5,6 % MgO (80 g/l water soluble magnesium oxide)

Mode of action and advantages:

- Water soluble phosphate for foliar application - directly available to plants.
- With magnesium and manganes for arable crops.
- For quick nutrient supply in times of high demand.

Recommendations for use and application rates:

General:	PhytoGreen®-HiPhos has to be applied as foliar fertiliser in at least 200 l Water/ha.
Potatoes:	To encourage tuber numbers, apply 10 l/ha in at least 200 litres of water at tuber initiation. To encourage bulking, apply 5 l/ha in at least 200 litres of water at the onset of bulking and repeat at least once during bulking. The timing of the 2nd application should be based on tissue analysis and should not be less than 10 days after the 1st.
Cereals:	Apply 5 l/ha in at least 200 litres of water at tillering and repeat at 10 - 14 day intervals as required. An application between ear emergence and the end of flowering may also be beneficial.
Oilseed rape:	Apply 2.5 - 5.0 l/ha in at least 200 litres of water. Apply in the autumn when the crop has 6 - 8 leaves and repeat in the spring at early stem extension.
Peas and beans:	Apply 5 l/ha in at least 200 litres of water when the crop is 10 - 15 cm high and repeat after 10 - 14 days.
Sugar beet:	Apply 5 l/ha in at least 200 litres of water at the 4 - 6 leaf stage. Repeat at 10 - 14 day
Corn:	Apply 2.5 - 5.0 l/ha in at least 200 litres of water at the 4 - 8 leaf stage. Repeat at 10 - 14 day intervals if required.



Corn with P-deficiency (purple leaves)

Technical details:

Density: 1,44 kg/l
pH = 2

Miscibility: Do not mix with the fungicide Corbel, with fertilisers containing boron or with alkaline products.

Pack sizes:
10 ltr. · 200 ltr. · 1000 ltr.

PHYTO solution
... Plant nutrition with system